

A Geostatistical Analysis of Water Losses in an Urban Utility using the R and Tinn-R Open-Source Statistical Computing Environments

F.Kizito^a, G. Ngirane-Katashaya^b & R. Thunvik^a

^aRoyal Institute of Technology (KTH), Sweden

^bMakerere University, Uganda



THE STUDY AREA:




THE STUDY AREA:

NINE
BRANCHES

3 - 7
TERRITORIES
PER BRANCH



The Kampala Water Supply Service Area



Frank Kizito, 2008

PROBLEM ISSUES:

Problem 1: Water Supply Intermittence




HAPHAZARD PIPE INFRASTRUCTURE
(Photo: KW Archives)


PROBLEM ISSUES:

Problem 2: High Non-Revenue Water (NRW) levels

| | | | | |
|---------------------|------------------------|---------------------------------|--------------------------------|-------------------|
| SYSTEM INPUT VOLUME | Authorised Consumption | Billed Authorised Consumption | Billed Metered Consumption | REVENUE WATER |
| | | | Billed Unmetered Consumption | |
| | | Unbilled Authorised Consumption | Unbilled Metered Consumption | NON-REVENUE WATER |
| | | | Unbilled Unmetered Consumption | |
| | Water Losses | Apparent (Commercial) Losses | Unauthorised use (Theft) | |
| | | | Metering Inaccuracies | |
| | | Real (Physical) Losses | Leaks | |
| | | | Bursts | |



NRW Performance Trends (July-December 2006)




LEAKS AND BURSTS
(Photo: KW Archives)


PROBLEM ISSUES:

Problem 3: Rising Production Costs vs Stagnant Collections

Ggaba Water Production 1992 -2007




Revenue Collections (Ushs 000' excl VAT)



GABA III WATER WORKS
(Photo: Roger Thunvik)



PROBLEM ISSUES:

Problem 4:
Increase in
disconnections due
to non-payment



A TYPICAL BRANCH OFFICE
(Photo: KW Archives)

STUDY OBJECTIVE:



To investigate the possible causes of declining water sales in a selected Branch, using geostatistical analysis.

TOOLS USED:



R version 2.6.1 (2007-11-26)
Copyright (C) 2007 The R Foundation for Statistical Computing



Tinn-R - GUI/Editor for R language and environment

Version 1.19.4.7

<https://sourceforge.net/projects/tinn-r>

<http://www.sciviews.org/Tinn-R>

Copyright 2001-2008

Under the GNU General Public License - GPL

R: A statistical computing environment, developed under the Free Software Foundation's GNU project.


(<http://www.r-project.org/>;
<http://www.gnu.org/>)

Tinn R: A free, simple code editor for R.
(<http://sourceforge.net/projects/tinn-r/>;
<http://www.sciviews.org/Tinn-R/>)

GEOSTATISTICAL ANALYSES:

(1) Selection of calibration and validation datasets

Spatial distribution of calibration and validation datasets



```
# Random selection of calibration and validation datasets
cal.id <- sample(1:dim(d200312.all@data)[1], 4592)
sort(cal.id)
d200312.cal <- d200312.all[cal.id, ]
str(d200312.cal)
summary(d200312.cal)
d200312.val <-
d200312.all[!is.element(1:dim(d200312.all@data)[1], cal.id),
]
str(d200312.val)
summary(d200312.val)


# Save calibration and validation datasets
save(d200312.all, d200312.cal, d200312.val, file =
"Project.RData")

# Spatial distribution of calibration and validation datasets
plot(coordinates(d200312.cal), col = "red", asp = 1, pch =
20, cex = 0.8, xlab = "UTM E (m)", ylab = "UTM N (m)", main =
and validation datasets", sub = selection))
points(coordinates(d200312.val), col = "blue", pch = 20, cex
= 0.8)
grid(lty = 500, col = "black")
```


GEOSTATISTICAL ANALYSES:

(2) Exploratory Data Analysis


Histogram of Monthly Water Sales Volumes (Calibration Dataset)




Histogram of Monthly Water Sales Volumes (Validation Dataset)



Classified Boxplot of Water Sales (calibration dataset)



Classified Boxplot of Water Sales (validation dataset)



Univariate exploratory graphics

#(a) Histograms

```
hist(d200312.cal$V200312, breaks = seq(0, 700, by = 20), col = "red", border = "darkblue", main = "Dataset", xlab = "Volume (m3)")
```

```
rug(d200312.cal$V200312)
```

```
hist(d200312.val$V200312, breaks = seq(0, 900, by = 20), col = "blue", border = "black", main = "Histogram of Monthly Water Sales Volumes (Validation Dataset)", xlab = "Volume (m3)")
```

```
rug(d200312.val$V200312)
```

#(b) Classified Boxplot


```
par(las=1)
```

```
par(cex.axis=0.6)
```

```
boxplot(d200312.cal$V200312 ~ d200312.cal$CCat, horizontal = T, outline = T, log="x", pars = list(boxwex = 0.4, staplewex = 0.3, outwex = 0.3), col = c("hotpink", "skyblue", "azure", "turquoise", "yellow", "tan"), main = "Classified Boxplot of Water Sales (calibration dataset)", xlab = "Volume (m3)",)
```

GEOSTATISTICAL ANALYSES:

(3) Investigation of spatial structure



Postplot of consumption categories

```
print(spplot(d200312.cal, z = "CCat", col.regions = bpy.colors(6), main = "Postplot of customer locations", sub = "Colour ramp, by consumption category", key.space = "inside", cex = 0.8))
```

Postplots of water sales values

##(a) Colour ramp, by deciles of log10 volume


```
print(spplot(d200312.cal, z = "log10V200312", col.regions = topo.colors(64), main = "Postplot of water sales volumes", sub = "log10 volume", key.space = "inside", cuts = quantile(d200312.cal$log10V200312, seq(0, 1, by = 0.1)), cex = 0.8))
```

##(b) Symbol size representing log10 volume

```
print(bubble(d200312.cal, "V200312", main = "Postplot of water sales volumes", sub = "Volume, symbol size proportional to value", col = "red", scales = list(draw = TRUE), xlab = "UTM E (m)", ylab = "UTM N (m)", fill = T, aspect = "iso", do.sqrt = T, key.entries = quantile(d200312.cal$V200312, seq(0, 1, by = 0.1))))
```

GEOSTATISTICAL (4) Variogram modelling

ANALYSES:



```
# Empirical variogram of logarithm of water sales volumes
```

```
v <- variogram(log10(V200312) ~ 1, loc = d200312.cal)
```

```
print (v)
```

```
print(plot(v, plot.numbers = T, main = "Empirical variogram, log10(Water sales volumes)", xlab = "separation distance, m", col = "darkblue", pch = 20))
```

```
# Fitting variogram model
```

```
vm <- vgm(0.04, "Sph", 1200, 0.13)
```

```
vmf <- fit.variogram(v, vm)
```

```
(vm <- vgm(0.005, "Sph", 5000, add.to = vmf))
```

```
(vmf2 <- fit.variogram(v, vm))
```

```
# Model validation using validation dataset
```

```
##(a) Add log10 volume field to validation dataset
```

```
d200312.val$log10V200312 <- log10(d200312.val$V200312)
```

```
##(b) Predict log10 water sales volumes at validation points
```

```
k.val <- krige(log10(V200312) ~ 1, loc = d200312.cal, newdata = d200312.val, model = vmf2)
```

```
summary(k.val)
```

```
##(c) Compare predictions to actual values
```

```
resid <- d200312.val$log10V200312 - k.val$var1.pred
```



```
summary(resid)
```

```
hist(resid, main = "Actual - Predicted, log10 water sales volumes, by OK", xlab = "log10 Volume", col = "lightblue")
```

```
resid <- as.data.frame(resid)
```


```
coordinates(resid) <- coordinates(d200312.val)
```

```
bubble(resid, maxsize= 2, main = "Residuals: Actual - Predicted, by OK", sub = "log10 Volume")
```



GEOSTATISTICAL ANALYSES:

(5) Water sales kriging predictions



Raster map of predictions

#(a) Selection of grid spacing

```
vc <- variogram(log10(V200312) ~ 1, loc = d200312.cal, cloud = T)
```

```
str(vc)
```

```
min(vc$dist)
```

```
(min.x <- floor(bbox(d200312.cal)["XCoord", "min"] - 10))
```

```
(min.y <- floor(bbox(d200312.cal)["YCoord", "min"] - 10))
```

```
(max.x <- ceiling(bbox(d200312.cal)["XCoord", "max"] + 10))
```

```
(max.y <- ceiling(bbox(d200312.cal)["YCoord", "max"] + 10))
```

```
(cells.x <- (max.x - min.x)/50)
```

```
(cells.y <- (max.y - min.y)/50)
```

```
cells.x * cells.y
```

#(b) Creation of grid

```
d200312.grid <- SpatialGrid(GridTopology(c(min.x + 25, min.y + 25), c(50, 50),
```

```
c(cells.x, cells.y)))
```

```
str(d200312.grid)
```

```
d200312.grid <- SpatialPoints(d200312.grid)
```

```
gridded(d200312.grid) <- T
```

```
str(d200312.grid)
```

#(c) Prediction on grid

```
k.grid <- krige(log10(V200312) ~ 1, loc = d200312.cal, newdata = d200312.grid, model = vmf2)
```

```
summary(k.grid)
```


#(d) Plotting gridded predictions

```
print(spplot(k.grid, zcol = "var1.pred", pretty = T, contour = T, col.regions = bpy.colors(64), main = "Predicted values, log10 water sales volumes", xlab = "UTM E", ylab = "UTM N", scales = list(draw = T)))
```

GEOSTATISTICAL ANALYSES:

(6) Difference map, highlighting areas with declining water sales

Difference in log10 water sales volumes, 2003 and 2007



```
# Compare 2003 predicted values and 2007 predicted values  
and generate a difference map
```

```
str(k.grid)
```

```
str(kComp.grid)
```

```
diff <- (k.grid@data$var1.pred - kComp.grid@data$var1.pred)
```

```
str(diff)
```

```
diff <- as.data.frame(diff)
```

```
str(diff)
```


```
coordinates(diff) <- coordinates(d200312.grid)
```

```
str(diff)
```

```
print(spplot(diff, zcol = "diff", cuts = quantile(diff$diff, seq(0, 1, by  
= 0.1)), pretty = T, contour = T, col.regions = topo.colors(64),  
main = "Difference in log10 water sales volumes, 2003 and  
2007", sub = log10Volume(2003) - log10 Volume(2007)", xlab =  
"UTM E", ylab = "UTM N", scales = list(draw = T), key.space =  
"right"))
```

GEOSTATISTICAL (7) Overlays of reported field anomalies

ANALYSES:



```
# Load datasets of anomalies
```

```
NoWater <- read.table("NoWater.dat", header = T)
```

```
coordinates(NoWater) <- c("XCoord", "YCoord")
```

```
AgedMeter <- read.table("AgedMeter.dat", header = T)
```

```
coordinates(AgedMeter) <- c("XCoord", "YCoord")
```

```
Disconnected <- read.table("Disconnected.dat", header = T)
```

```
coordinates(Disconnected) <- c("XCoord", "YCoord")
```

```
Estimated <- read.table("Estimated.dat", header = T)
```

```
coordinates(Estimated) <- c("XCoord", "YCoord")
```

```
# Postplots of anomalies, superimposed on difference map
```

```
##(a) No water cases
```

```
plot(coordinates(diff), cex = 1.5*(diff$diff)/max(diff$diff), pch = 20, col = "skyblue", main = "Difference map, with 'no water' cases", sub = high differences in water sales; red dot = 'no water' case", xlab = "UTM E", ylab = "UTM N")
```

```
points(coordinates(NoWater), col = "red", pch = 20, cex = 1.3)
```

```
##(b) Aged meters
```

```
plot(coordinates(diff), cex = 1.5*(diff$diff)/max(diff$diff), pch = 20, col = "skyblue", main = "Difference map, with aged meter locations", sub = areas = high differences in water sales; red dot = aged meter", xlab = "UTM E", ylab = "UTM N")
```

```
points(coordinates(AgedMeter), col = "red", pch = 20, cex = 1.3)
```

RESULTS AND CONCLUSIONS:

- ▶ Out of eight factors investigated, “intermittent supply” cases correlated most with areas of declining water sales.
- ▶ Aged meters were also seen to have a contributing effect to low water sales.
- ▶ Disconnected and inactive accounts were numerous and widespread throughout the study area.
- ▶ Leaks and illegal cases seemed to have been too few to impact significantly on water sales.

ACKNOWLEDGMENTS

This study was carried out within the framework of the Sida/SAREC-funded project:

“Sustainable Technological Development in the Lake Victoria Basin”





Thank You!

Asanteni!

TACK!!

